



Agromet Advisory Bulletin

Date : 12-12-2023

Weather Forecast of **UMRED** Block in **NAGPUR**(Maharashtra) Issued On :2023-12-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-12-13	2023-12-14	2023-12-15	2023-12-16	2023-12-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	28.2	28.5	28.9	29.1	29.4
Tmin(°C)	15.6	15.2	14.4	14.5	14.7
RH-I(%)	80	76	73	71	72
RH-II(%)	62	59	56	54	56
Wind Speed(kmph)	4	5	5	5	5
Wind Direction(Degree)	58	49	55	47	48
Cloud Cover(Octa)	6	6	2	1	1

Weather Summary/Alert:

• As per the blocklevel value added forecast given by, IMD, RMC, Nagpur, sky will be clear to partly cloudy during next five days i.e.,13th to 17th, December, 2023. • Weather is very likely to be dry on 13th, 14th, 15th, 16th and 17th, December, 2023. • No large change in maximum and minimum temperature during next 5 days over the Vidarbha.

General Advisory:

• Harvested paddy crop which has been exposed to rains of previous week should be sun dried and threshed keeping in view dry weather forecast. • Considering the dry weather forecast, it is advised to continue the agrochemical spraying operations, intercultural operations and fertilizer application in standing crops during next 5 days. • It is advised to continue the harvesting and threshing of matured paddy crop. • It is advised to continue to picking of burst cotton and keep the picked cotton picking and variety wise. Use cotton bags instead of gunny or plastic bags for picking and storing of cotton to avoid contamination of the lint.

SMS Advisory:

• Considering the dry weather forecast, it is advised to continue the agrochemical spraying operations, intercultural operations and fertilizer application in standing crops during next 5 days.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
PIGEON PEA (RED GRAM/ARHAR)	• Due to rainfall and continuous cloudy weather during previous week, to avoid the flower dropping in arhar spraying should be done with NAA 4 ml + 10 ml boron in 15 liters of water during clear weather condition. • Install pheromone traps at a distance of 50 m @ 5 / ha for monitoring of Helicoverpa. If the incidence of pod borer is noticed on pigeon pea crop and crossing economic threshold level, spraying should be done of any of recommended insecticide viz., Chlorantraniliprole 18.50 % SC @ 150 ml in 500 to 750 litres of water per hectare or Emamectin benzoate 05 % SG @ 220 gram in 500 to 750 litres of water per hectare or Ethion 50 % EC @ 1000-1500 ml in 500-1000 litres of water per hectare or Flubendiamide 20 % WG @ 250 gram in 500 litres of water per hectare or Flubendiamide 39.35 % m/m SC @ 100 ml in 500 litres of water per hectare or Lambda-cyhalothrin 05 % EC @ 400-500 ml/ha in 400-600 liters of water per hectare or Quinalphos 25 % EC @ 1400 ml/ha in 500-1000 liters of water per hectare or Chlorantraniliprole 09.30 % + Lambda-cyhalothrin 04.60 % ZC 200 ml/ha in 500 litres of water per hectare.
BENGAL GRAM/ CHICK PEA	• Due to rainfall and continuous cloudy weather prevail during previous week, to prevent the occurrence of wilt disease in gram crop, excess water accumulated in the field should be drained immediately and drenching of Trichoderma biological fungicide mixed with 40 grams per 10 liters of water should be sprayed or drenched or thiophanate methyl 70 WP 25 grams mix with 10 liters of water should be sprayed. • For integrated management of Gram Pod Borer 20 bird perches per hectare should be installed up in the field. For control of gram pod borer install pheromone traps (Hexalure) two per acre or five per hectare. If eight to ten moths are found in the trap for three consecutive days, recommended management measures should be taken. In case of pest infestation by observing the crop or when the crop is in 40 to 50 % flowering stage, preference should be given to botanicals or biological pesticides. For this, first spraying should be taken of neem extract 5% or Azadiractin 300 ppm 50 ml per 10 liters of water with Knapsack sprayer pump. If spraying should be done with a power sprayer, apply three times the amount of pesticide. If Gram Pod Borer has reached the ETL, spraying should be done with Quinalphos 25 % EC 20 ml or Emamectin benzoate 5 % SG 3 gm or Flubendiamide 20 % WG 5 gm or Chlorantraniliprole 18.5 SC 2.5 ml per 10 liters of water. At Wafsa condition hoeing

Crop(Varieties)	Crop Specific Advisory
	and weeding must be carried out twice; first at 20 and second at 30 days old plants to ensure weed free environment during the critical period of crop weed competition for better water and nutrient use efficiency.
MUSTARD	<ul style="list-style-type: none"> Cloudy weather and lowering night temperature favours aphid incidence in mustard. For control of aphid incidence undertake spray of Thiometon 25 EC 8 ml per 10 lit of water or Dimethoate 30 EC 10 ml per 10 lit of water. Undertake need based irrigation for higher productivity in mustard. In case of availability of one irrigation schedule it at flowering stage, for availability of two irrigations schedule it at 30 days and flowering stage, for availability of three irrigations schedule it at 25 to 30 days interval.
CITRUS	<ul style="list-style-type: none"> If Orange/Mosmbi/Lemon Ambia are left under stress, the unseasonal rains break the stress, spray Cycocel (chlormequat chloride 50% SL)* 02 ml per liter of water to keep the orchards under stress. Orchards which were not originally stressed should be sprayed with chlormequat chloride 50% SL at 04 ml per liter of water. To avoid the fruit dropping and to increase the size of fruit of orange and Sweet orange of Mrig bahar and Acid lime of Hastha bahar, spraying should be done of any one of NAA (01 gm) or 2,4-D (1.5 gm) or gibberellic acid (1.5 gm) along with 6 BA (01 gm) + 13:00:45 (01 kg) + propiconazole 25% EC* 100 ml mixed with 100 liters of water and a sticker should be used as needed. In Ambia bahar and Mrig bahar orchards due to continuous cloudy weather and injury after rains the entire plant should be sprayed with copper oxychloride 50 WP 30 g mixed with 10 liters of water. Due to rain, there is a large incidence of citrus canker disease on acid lime, so for protection, copper oxychloride 30 gm + streptomycin 1 gm mixed with 10 liters of water should be sprayed. Special Notice: * Please note that some of the chemicals and biocides mentioned in this do not have label claims.
RICE	<ul style="list-style-type: none"> It is advised to continue the harvesting and threshing of matured paddy crop. Harvested paddy crop which has been exposed to rains of previous week should be sun dried and threshed keeping in view dry weather forecast.
WHEAT	<ul style="list-style-type: none"> For late sowing of irrigated wheat 150 kg seed per hectare should be used for sowing. For coarse seed varieties like HD 2189 or Purna of wheat crop, 125 kg seed per hectare should be used. Apply 80 kg N, 40 kg P and 40 kg K per hectare for late sowing wheat. For both irrigated and late sowing, apply half dose of nitrogen as well as full phosphorus and potash at the time of sowing and the remaining half dose at the time of first irrigation (18 to 20 days after sowing).
COTTON	<ul style="list-style-type: none"> It is advice that in areas where the cotton crop is ready for picking give priority for clean picking of cotton. Keep the picked cotton, variety wise. Use cotton bags instead of gunny or plastic bags for picking and storing of cotton to avoid contamination of the lint. If pink bollworm incidence crosses ETL, (i.e. 5-10% of green boll infestation or more than 8 moths per trap per night consecutive 3 days), spray Cypermethrin 10% EC @10-15ml Or Cypermethrin 25% EC @ 46ml Or Lambda cyhalothrin 5%EC @10ml Or Deltamethrin 2.8EC @10ml Or Fenprothrin 10 % EC@15-20ml or Fenvalerate 20EC @ 10ml Or Alphacypermethrin 10% EC @ 6ml/10 litres of water. To manage grey mildew and leaf spots, undertake spray of Carbendazim 12%+ Mancozeb 63% WP@30 g or Kresoxim methyl 44.3 SC @ 10 ml or Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC @ 10 ml in 10 litres of water. Apply foliar spray of copper oxychloride 50 WP/WG@ 25-30 g/10 litres of water to manage bacterial blight in cotton. Collect and destroy diseased bolls and crop debris from the fields.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
LINSEED	<ul style="list-style-type: none"> At Wafsa condition, carry out first hoeing after 25 days after sowing. Hand weeding should be done keep in the of weed intensity.
CHILLI	<ul style="list-style-type: none"> Due to alternative low and high temperature coupled with high humidity, if the powdery mildew disease is noticed on chilli crop, spraying should be done of any of following fungicide, Hexaconazole 75 % WG @ 66.7 gram or Tebuconazole 25% WG @ 500-750 gram or Azoxystrobin 8.3 % + Mancozeb 66.7 % WG @ 1500 gram or Boscalid 25.2% + Pyraclostrobin 12.8 % WG @ 600 gram or Carbendazim 12 % + Mancozeb 63 % WP @ 750 gram or Kresoxim-Methyl 15 % + Chlorothalonil 56 % WG @ 1000 gram or Tebuconazole 10 % WP + Sulphur 65 % WG @ 1250 gram or Tebuconazole 50 % + Trifloxystrobin 25% WG @ 250 gram mixed with in 500 litre of water per acre for management of disease.
BRINJAL	<ul style="list-style-type: none"> For management of brinjal fruit and shoot borer Spray Bacillus thuringiensis formulation (1 ml/L) at weekly interval Release Trichogramma chilonis @ 2,50,000/ha (50,000/release-5 times at weekly intervals, starting from flowering). Install sex pheromone traps 10/acres. Spray Chlorantraniliprole 18.5 SC at 0.3 ml/L once in 15 days depending upon the pest population.
TOMATO	<ul style="list-style-type: none"> As a precautionary measure, for management of Early Blight disease of tomato spraying should be done with Copper oxychloride (3.0 g/l) or Copper hydroxide (2.0 g/l) one or two days before transplanting. Spray Copper oxychloride (3.0 g/l) or Mancozeb (2.0 g/l) or Chlorothalonil (2.0 g/l) or Propineb (2.0 g/l) or Metiram (2.0 g/l) or Pyraclostrobin + Metiram (2.0 g/l) or Tebuconazole 50 % + Trifloxystrobin 25 % w/w (0.6 ml/l) at 10-15 days interval or as and when required in the main field. If the symptoms of Late Blight of tomato was noticed due to the previous cloudy weather, spraying should be done with Mancozeb (2.0 g/l) or Copper oxychloride (3.0 g/l) or Bordeaux mixture (1%) or Copper hydroxide (2.0 g/l) or Fosetyl-Al (2.0 g/l) or Dimethomorph (2.0 g/l) combination of Pyraclostrobin + Metiram (2.0 g/l) or

Horticulture(Varieties)	Horticulture Specific Advisory
	Mefenoxam + Copper hydroxide (2.0 g/l) or Metalaxyl 8% + Mancozeb. 64% (2.0 g/l) during clear weather condition.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
BUFFALO	<ul style="list-style-type: none"> To protect livestock from cold weather, keep the animals indoor during night hours. Provide suitable bedding like paddy straw, wheat straw, saw dust etc. of 4-6-inch thickness to protect the animals from cold. The floor of the animal shed should be kept clean and dry. Provide clean and potable drinking water to animals round the clock. Clean the water trough (once in a week) regularly in livestock. Provide mineral mixture @ 30-40 g/day to cattle for improving the milk production and reproduction efficiency.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	<ul style="list-style-type: none"> It should be noted that the validity of the dates mentioned in the Weather Based Advisory Bulletins will be valid up to 08:30 AM of the mentioned date. *Based on a research trial that was not included in the CIBRC label claim. # Under label claim.
GENERAL ADVICE	<ul style="list-style-type: none"> Nursery brinjal, tomato, cauliflower and cauliflower seedlings should be planted in the field at the age of 4 to 6 weeks. Fenugreek, spinach, cilantro, radish and carrot should be planted in stages.