



**Gramin Krishi Mausam Sewa**  
**Agromet Advisory Bulletin for NAGPUR District**  
 Central Institute for Cotton Research,  
 Nagpur



## Agromet Advisory Bulletin

Date : 2025-08-19

Weather Forecast of District NAGPUR (Maharashtra) Issued On : 2025-08-19 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-08-20	2025-08-21	2025-08-22	2025-08-23	2025-08-24
Rainfall(mm)	124.4	12.5	11.0	8.3	10.7
Tmax(°C)	29.4	28.6	29.1	28.9	29.1
Tmin(°C)	24.2	24.0	23.9	23.5	23.4
RH-I(%)	98	95	91	91	90
RH-II(%)	84	73	75	75	76
Wind Speed(kmph)	3	10	14	13	15
Wind Direction(Degree)	310	231	238	247	247
Cloud Cover(Octa)	8	7	7	7	8
Warning	Very Heavy Rain; Thunderstorms & Lightning, Squall etc	Thunderstorms & Lightning, Squall etc	No Warning	No Warning	Thunderstorms & Lightning, Squall etc

### Forecast Summary:

• In Nagpur district, as per the district level value added forecast given by, IMD, RMC, Nagpur, sky will be mostly cloudy during next five days i.e. 20 to 24 August, 2025. • Light to moderate rainfall is very likely to occur at most places on 20 and 21, August, 2025. • Light to moderate rainfall is very likely to occur at many places on 22 August, 2025. • Light to moderate rainfall is very likely to occur at few places on 23 and 24 August, 2025. • There will be fall in maximum temperature by 2 – 3 °C in the next 24 hours, thereafter no large change in maximum temperature over Vidarbha region. • As per the Sub Division wise Extended Range Rainfall Forecast, above normal rainfall, below normal maximum temperature and normal minimum temperature is likely to be occurring over Vidarbha during 24 to 30, August, 2025.

### Weather Warnings (Valid Till 08:30 IST of the next day)

• Heavy to very heavy rainfall with thunderstorm accompanied with lightning likely to occur at one or two places on 20 August, 2025. • Thunderstorm accompanied with lightning likely to occur at one or two places on 21 and 24 August, 2025.

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Lightning strikes accompanied by thunder can cause injury or death to livestock, farmers or farm workers. Farmers and farm workers engaged in paddy planting or weeding seasonal crops can be injured or killed by lightning. Animals can also be swept away by flowing water. If rainwater remains stagnant in crop/vegetable and orchard areas for a long time, crop growth can be stunted and diseases such as blight, root rot etc. can spread in the crop.

**General Advisory:**

• Agrochemicals spraying, fertilizer application, intercultural operations and transplanting operations should be postponed for next 3 days on accounts of forecasted widespread rainfall along with thunderstorm, lightning, gusty winds and heavy rainfall. • During very essential farm operations, farmers and farm workers should avoid working together in large numbers while transplanting paddy or other plants, and maintain maximum distance between two farm workers while performing essential agricultural work. During such work, use the mobile app Damini Lightning Alert to monitor lightning locations. If you are in a lightning-prone area, immediately seek shelter in a safe place. • Spraying of essential agricultural chemicals should be done during local clear and calm weather conditions after the current spell rainfall. • It is advised to support the fruit orchards and vegetables with the help of Bamboos to avoid the fruit dropping. • Open the furrows, In orchards, vegetable crops, as well as crops like soybean, cotton, tur, to prevent it from rainwater stagnated. Excess water accumulated in the crop area should be drained. • Farmers, farm laborers should not rush to cross the road if water is flowing from the river and nala and also take care that their other animals do not pass through the flowing water. • Cows, buffaloes, goats, sheep and other domestic animals should be avoided to graze in the open spaces on days when thunders are predicted. Animals should be kept away from open water sources, rivers or lakes and away from tractors and other metal implements. Sufficient fodder and water should be arranged for the animals in the manger. Farmers and farm labourers should take care of themselves and livestock keeping in mind the forecast of lightning. Sheltering under trees should be strictly avoided and also livestock should not shelter under trees.

**SMS Advisory:**

• Install pheromone traps @ 5 per hectare for monitoring moth activity of pink bollworm.

**Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory
SOYABEAN	<p>• It is necessary to remove the accumulated water in the field from the field in the direction of the slope. It is necessary to spray 00:52:34 (Nitrogen: Phosphorus Potassium) soluble fertilizer at a rate of 1% (one kg of soluble fertilizer in 100 liters of water). Along with the main nutrients, micronutrients such as ferrous, zinc, magnesium should be sprayed. Micronutrient deficiency is felt in the soybean crop at the stage of flowering and pod setting, in such a situation, spraying with micronutrient grade-2 (2.5 to 3 ml per liter of water) helps in filling the pods well. To maintain genetic purity in the seed-producing field, plants of different varieties should be removed from your field based on flower color and leaf/leaf/pod color. • There is a possibility of infestation of tobacco leaf-eating caterpillars, sedge caterpillars and green camel caterpillars on the soybean crop. For this, it is necessary to spray insecticides such as Emamectin Benzoate 5% SG (200 g/ha) plus Thiomethoxam 25% WG (100 g/ha) or Indoxacarb 14.5% SC 500 ml per hectare or Emamectin Benzoate 5% SG 200 g/ha or Thiomethoxam 25% WG 100 g/ha or Chlorethrinilliprol 18.5% W/W SC 150 ml per hectare. • Rhizoctonia aerial blight on soybean crops is mainly seen in moist and humid weather, if there is heavy rainfall, riverbank land, waterlogged fields, the disease is seen in those places. Preventive measures are necessary. For this, any one of the following fungicides should be sprayed. Pyroclostrobin 20% WG @ 375-500 gm per hectare (1 gm per litre water) or Fluxapyroxad Pyroclostrobin @ 300 ml per hectare (0.60 ml per litre water) or Pyroclostrobin Epoxyconazole 750 ml per hectare (1.5 ml per litre water). • Spray any one of the following fungicides for the management of scab disease in soybean crop. Tebuconazole 10% + Sulphur 65% WG 1.25 kg/ha (2.5 gm per litre water) or Carbendazim 12% + Mancozeb 63% WP (2 gm/litre water) or Tebuconazole 29.9 EC 625 ml/ha (1.25 ml/litre water). • At present, there is moisture in the soil, but if there is a break in rains and the temperature increases during the day, the incidence of charcoal rot may increase. As soon as the symptoms of this disease appear, a protective irrigation should be done. Trichoderma, a biological fungicide (5-10 grams per liter of water) should be applied in patches. This will prevent the disease from growing in the field and the severity of the disease can be reduced. As soon as the symptoms of this disease appear, spray Carbendazim 12% Mancozeb 63% WP (2.5 grams/liter of water) or Fluxapyroxad +</p>

Crop (Stage)	Crop Specific Advisory
	<p>Pyroclostrobin (1 gram per liter of water) and ensure that the fungicide reaches the lower part of the plants. • In the present situation, the incidence of yellow mosaic disease is more or less visible on soybean crops. This disease is mainly spread by whiteflies. Yellow sticky traps should be placed at the rate of 50 per hectare. If the incidence of yellow moth is low, the infected plants in the field should be removed and buried in the ground. To control the sucking insects, the following inter-flow insecticides should be sprayed. Thiomethoxam Lambda-Cyhalothrin 125 ml per hectare (0.25 ml per liter of water) or Beta-Cyhalothrin Imidacloprid 350 ml per hectare (0.7 ml per liter of water) or Acetamiprid Bifenthrin 250 g per hectare (0.5 g per liter of water) should be sprayed.</p>
PIGEON PEA (RED GRAM/ARHAR)	<p>• Nipping (Removing the apical bud) is beneficial for increasing tur (pigeon pea) production: - The first nipping should be done about 40 to 45 days after sowing and the second nipping should be done after 60 to 65 days so that more shoots and branches are produced. Nipping should be done by hand to make them two to three inches deep. After nipping, applying phosphorus fertilizer also helps in the establishment of flowers and pods, which increases the yield.</p>
RICE	<p>• Spraying of essential agricultural chemicals should be done during local clear and calm weather conditions after the current spell rainfall. • Marigold and Cowpea crops should be planted on paddy bunds for breeding of beneficial insects. • Nursery: - For management of Stem borer and Gall midge apply Carbofuron 3% granular 25 kg per hectare (Carbofuron 3% granular 250 gm per 100 square meter nursery) should be applied in the nursery 5 days before transplanting. • Before transplanting roots of plants are treated with Chlorpyrifos 20% @10 ml solution. By soaking in the prepared solution for 12 hours per 10 liters of water and then transplant. • Paddy nursery should be kept free from weeds. • Marigold should be planted as a trap crop on paddy bunds. • The tops of the seedlings should be cut off and stored in bamboo baskets so that the eggs of the Stem borer on the seedlings are destroyed and parasitic insects come out of them. • Toxic gooseberry leaves at the rate of 1.5 tons per hectare should be incorporated in the field during monsoon, it also reduces the infestation of Stem borer and Gall midge and Leafhoppers. • Seedlings should be planted in a strip (Alleyways) system at the recommended spacing (20 x 15 or 20 x 20 cm) to manage the Leafhoppers. (Leave 30 cm space after 10 lines or 2 meter) • Chemical Fertilizers: - Agriculture is profitable if the soil is tested and proper quantities of chemical fertilizers are used. Generally, apply 100 kg Nitrogen, 50 kg Phosphorus and 50 kg Potash per hectare. Apply whole phosphorus and potash and half the amount of nitrogen in the mud (Puddled field) and the remaining half nitrogen in two equal installments (usually tillering stage which comes at 30 days and panicle initiation stage which comes about 60 days). • Pre-emergence: - Pendimethalin 30 % E. C. @50 ml in 4 to 7 days after transplanting or Pretilachlor 50 % E. C. @ 20 to 30 ml in 4 to 7 days after transplanting or Pyrazosulfuron ethyl 10% w.p. @ 2 to 3 grams mixed in 10 liters of water should be sprayed in 4 to 7 days after transplanting.</p>
COTTON	<p>• Spraying of essential agricultural chemicals should be done during local clear and calm weather conditions after the current spell rainfall. • Root rot disease to seedling plants. Diseased plants may be in patches. Spot/line application of early symptomatic plants with 1 kg Trichoderma viride/T. harzianum WP or liquid formulation mixed in well decomposed 50 kg FYM or compost for application in 1 acre area. OR Spot drenching with Carbendazim 50 WP @ 12 g mixed in 10 litres of water only, solution volume just sufficient to wet root zone. • Monitoring of pink bollworm using pheromone traps may be initiated 45 days after sowing. Install pheromone traps @ 5 per hectare for monitoring moth activity of pink bollworm.</p>

### Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
MANDARIN ORANGE	<p>• Spraying of essential agricultural chemicals should be done during local clear and calm weather conditions after the current spell rainfall. • Continuous drizzle and continuous cloudy weather favours fungal disease infection. Since the weather is conducive to the fungus that causes the</p>

Horticulture (Stage)	Horticulture Specific Advisory
	<p>diseases such as petiole drying, leaf spot, leaf drop and brown spot on fruits, one should know the symptoms of fruit drop in the garden and take appropriate measures. • Care should be taken that rainwater does not accumulate in the fruit orchards, the accumulated water should be drained immediately towards the slope of the land. • The fallen leaves and fruits should be disposed of. Leaf and decayed fruit should not be kept in orchards as these lead to the spread of pests and diseases. • As a preventative measure to prevent the spread of leaf drop and brown rot on fruits caused by Phytophthora fungus, spray the entire plant with Fosatil AL * 2.5 gm or Copper Oxychloride * 50 WP 3 gm per liter of water. After spraying with agrochemicals Trichoderma harzianum* plus Sudomonas fluorescens* 100 gms each mixed with 1 kg of FYM should be applied from the soil around the tree. • (*recommendation based on research, not a label claim) • For ambia bahar crop of Nagpur mandarin and sweet orange at fruit development stage, apply 50 gm of ferrous sulphate (FeSO<sub>4</sub>) and 50 gm of zinc sulphate (ZnSO<sub>4</sub>) and 5 kg cow dung or vermicompost along with application of remaining recommended dose of fertilizer. • In the root rot affected orchard, apply and spraying should be done of Mefenoxam MZ 2.5 gram per litre of water and 5-10 liters (depending on tree size) per tree or fosetyl aluminium 2.5 g/litre (twice at 40 days interval). All fruits on the affected plants should be removed and then sprayed with a solution of Mefenoxam MZ (2.5 g per liter of water) + Carbendazim (1 g per liter of water). • Carbendazim or thiophanate methyl should be sprayed twice at the rate of 1 gram per liter of water on plants affected by Colletotrichum and Botryodiplodia fungi. After 10 days after the first spraying, the second spraying should be repeated. Azoxystrobin + difenconazole fungicide 0.5ml per litre of water can be sprayed as an alternative. • Install methyl eugenol pheromone trap at the rate of 20-25 hectare control of for fruit fly. Bait (methyl eugenol) should be changed every 15-20 days. • For the management of fruit sap sucking moth at the time of colour development stage, spraying should be done of Neem oil 10 ml or petroleum spray oil (mineral oil) ml per litre of water 10-15 days interval, till the fruits are harvested. • • Brown rot on fruits and fruit rot management: First of all, the fallen leaves and fruits should be disposed of and should not be left in the field, otherwise it will help in increasing the severity of the disease and the infection will spread rapidly. Keep the soil clean. • • The water in the field should be drained from the slope of the garden because the area where water accumulates is more prone to Phytophthora fungus infection. • • As a preventive measure to prevent the infestation, spray the entire tree with Facitil A.L. 2.5 gm Kiwan Copper Oxychloride 50 WP 3 gm per liter of water. While spraying, spray the circumference of the tree as well so that if the fallen fruits have not been picked, the fungus on them will be destroyed and it will also help in destroying the active spores in the soil. For better results, do not mix any other similar fungicides/insecticides/soluble fertilizers with this medicine. • Metlaxyl M 3.3% + Chlorothlonil 33.1% SC (mixed fungicide) 2 ml per liter of water and trenching should be done (at least 5 liters of solution should be used per tree) • After 15 days of spraying chemical drugs, Trichoderma harzianum plus Pseudomonas fluorescens 100 g each per tree mixed with 1 kg of cow dung and applied through the soil around the tree. (Note: Based on research; Label claim recommendation) • • Reference: AICRP (Fruits), Dr. PDKV., Akola.</p>

### Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
COW	<p>During October, outbreaks of Fascioliasis, Lumpy Skin Disease (LSD), Haemorrhagic Septicaemia (HS), and Enterotoxaemia (ET) are likely in Maharashtra as per forewarning given by ICAR–National Institute of Veterinary Epidemiology and Disease Informatics (ICAR–NIVEDI), Bengaluru. To prevent losses, farmers are advised to follow these measures:</p> <ol style="list-style-type: none"> <li>Fascioliasis (Liver fluke disease) • Shed Management: <input type="checkbox"/> Avoid grazing in marshy/waterlogged pastures. <input type="checkbox"/> Keep sheds dry and ensure proper drainage around animal housing. • Feed &amp; Water Management: <input type="checkbox"/> Provide clean drinking water, preferably from borewell/tubewell (not stagnant ponds). <input type="checkbox"/> Avoid feeding fodder from swampy areas. • Prophylaxis: <input type="checkbox"/> Deworm animals</li> </ol>

Live Stock	Live Stock Specific Advisory
	<p>with flukicidal drugs (closantel, triclabendazole, rafoxanide) as per veterinarian's advice. • Ecto/Endoparasite Control: <input type="checkbox"/> Regular deworming schedule to break parasite cycle. 2. Lumpy Skin Disease (LSD) • Shed Management: <input type="checkbox"/> Install mosquito nets/screens; ensure proper ventilation. <input type="checkbox"/> Regular cleaning of sheds with lime/disinfectants. • Feed &amp; Water Management: <input type="checkbox"/> Provide balanced nutrition and mineral mixture to boost immunity. <input type="checkbox"/> Ensure adequate clean water supply. • Prophylaxis: <input type="checkbox"/> Vaccination with Goatpox/LSD vaccine (as per state veterinary department guidelines). • Ecto/Endoparasite Control: <input type="checkbox"/> Spray/smear insect repellents (cypermethrin, deltamethrin) in and around sheds. <input type="checkbox"/> Control vector population (mosquitoes, flies, ticks) o 3. Haemorrhagic Septicaemia (HS) • Shed Management: <input type="checkbox"/> Maintain hygiene; avoid overcrowding in sheds. <input type="checkbox"/> Protect animals from sudden weather changes (cold/rain). • Feed &amp; Water Management: <input type="checkbox"/> Provide adequate green fodder and concentrate feed. <input type="checkbox"/> Ensure fresh, uncontaminated water. • Prophylaxis: <input type="checkbox"/> Vaccinate susceptible cattle and buffaloes with HS vaccine before monsoon/post-monsoon. • Ecto/Endoparasite Control: <input type="checkbox"/> Deworm and maintain ectoparasite control to reduce stress load. 4. Enterotoxaemia (ET) – mostly in sheep &amp; goats • Shed Management: <input type="checkbox"/> Avoid sudden exposure to lush, high-protein pasture after dry periods. <input type="checkbox"/> Maintain dry, clean sheds to prevent stress. • Feed &amp; Water Management: <input type="checkbox"/> Avoid sudden change in diet. <input type="checkbox"/> Prevent overeating of concentrates; provide roughage before concentrates. • Prophylaxis: <input type="checkbox"/> Vaccination of sheep/goats with ET vaccine (Clostridial vaccine). • Ecto/Endoparasite Control: <input type="checkbox"/> Regular deworming to reduce gut load and improve immunity. General Advisory for Farmers: <input type="checkbox"/> Keep record of vaccination and deworming. <input type="checkbox"/> Report any unusual illness or sudden deaths immediately to nearest veterinary dispensary/KVK. <input type="checkbox"/> Ensure insurance coverage of animals wherever possible.</p>

### Likely impacts of weather warnings (General)

Lightning strikes accompanied by thunder can cause injury or death to livestock, farmers or farm workers. Farmers and farm workers engaged in paddy planting or weeding seasonal crops can be injured or killed by lightning. Animals can also be swept away by flowing water. If rainwater remains stagnant in crop/vegetable and orchard areas for a long time, crop growth can be stunted and diseases such as blight, root rot etc. can spread in the crop.

### Impact based advisories (General)

• Agrochemicals spraying, fertilizer application, intercultural operations and transplanting operations should be postponed for next 3 days on accounts of forecasted widespread rainfall along with thunderstorm, lightning, gusty winds and heavy rainfall. • During very essential farm operations, farmers and farm workers should avoid working together in large numbers while transplanting paddy or other plants, and maintain maximum distance between two farm workers while performing essential agricultural work. During such work, use the mobile app Damini Lightning Alert to monitor lightning locations. If you are in a lightning-prone area, immediately seek shelter in a safe place. • Spraying of essential agricultural chemicals should be done during local clear and calm weather conditions after the current spell rainfall. • It is advised to support the fruit orchards and vegetables with the help of Bamboos to avoid the fruit dropping. • Open the furrows, In orchards, vegetable crops, as w

**Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightning.**

**Mausam MobileApp link: <https://play.google.com/store/apps/details/>**

**Meghdoot MobileApp link: <https://play.google.com/store/apps/details>**

**Damini MobileApp link : <https://play.google.com/store/apps/details>**