



## Agromet Advisory Bulletin

Date : 28-09-2023

Weather Forecast of **PARSEONI** Block in **NAGPUR**(Maharashtra) Issued On :2023-09-28(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-29	2023-09-30	2023-10-01	2023-10-02	2023-10-03
<b>Rainfall</b>	11.1	13.6	16.0	17.8	22.7
<b>Tmax(°C)</b>	32.1	31.6	31.4	30.9	29.8
<b>Tmin(°C)</b>	23.3	23.0	23.4	23.6	23.8
<b>RH-I(%)</b>	83	85	89	91	88
<b>RH-II(%)</b>	63	65	69	71	68
<b>Wind Speed(kmph)</b>	5	5	5	5	5
<b>Wind Direction(Degree)</b>	209	194	299	226	330
<b>Cloud Cover(Octa)</b>	6	6	7	7	7

### Weather Summary/Alert:

- In the block, as per the block level value added forecast given by, IMD, RMC, Nagpur, sky will be partially to mostly cloudy during next five days i.e. 29th, September to 03rd, October, 2023. • Light to moderate rainfall likely to occur on 29th and 30th, September and 01st, 02nd and 03rd, October, 2023. • Thunderstorm with lightning likely to occur on 29th, 30th September and 01st, October, 2023. • As per the extended range forecast system (ERFS) rainfall will be above normal, maximum and minimum temperature will be normal during the period from 01st to 07th, October, 2023 over the Vidarbha region.

### General Advisory:

- Considering rainfall forecast it is advised to continue the agrochemical spraying operations and intercultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • Harvesting of matured green gram, black gram and early sown and early maturity soybean should be continue during next 1-2 days by taking in to account of local clear and calm weather condition. • Harvested crops should be stored at elevated places and covered with plastic sheets. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • Considering the thunderstorm and lightning forecast, it is advised to avoid letting cow, buffalo, goats, sheep, and other cattle's/animals graze in open spaces as there is a possibility of rain and lightning. Animals should be kept away from open water sources, rivers or lakes and tractors and metal. Cconsidering the thunderstorm and lightning forecast it is advised to carry out farm operations during early morning hours. Carry out farm operations by considering the local weather forecast and warnings. Take shelter at safe places, it is ensuring to keep maximum distance between two farm labours while working in the field. Avoid shelter near to water bodies (Lake, Well, River and Moist places) tractors and metal implements. Avoid taking shelter under a tree in any situation as the lightning attracts on tallest object.

### SMS Advisory:

- It is advised to continue the agrochemical spraying operations and intercultural operations during next 1-2 days by taking in to account of local clear and calm weather condition.

Crop(Varieties)	Crop Specific Advisory
<b>COTTON</b>	considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • If the incidence of internal boll rot/bacterial boll rot is noticed in cotton at squaring, flowering and boll development stage due to high humidity, cloudy weather and continuous rainfall during previous week, for management it is advised to collect and destroy the dried petals sticking to the developing bolls should be removed. Avoid indiscriminate use of nitrogenous fertilizers. Restrict excess vegetative growth of the cotton crop. Facilitate proper drainage in the field to avoid water logging in the field. Prophylactic sprays of Copper

Crop (Varieties)	ing in to account of local clear and calm weather condition.
Crop Specific Advisory COTTON	<p>considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • If the incidence of internal boll rot/bacterial boll rot is noticed in cotton at squaring, flowering and boll development stage due to high humidity, cloudy weather and continuous rainfall during previous week, it is advised to continue the agrochemical spraying operations and inter-cultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also drainage in the field to avoid water logging in the field. Prophylactic sprays of Copper Oxychloride 50 WP @25 g/10 L is suggested during early boll developmental stages at 15 days interval. Manage sucking pests with spray of recommended insecticides. • If the incidence Target leaf spot, Alternaria leaf spot, Myrothecium leaf spot, External fungal boll rot was noticed in cotton crop due to due to high humidity, cloudy weather and continuous rainfall during previous week, it is advised to take the prophylactic spray of Propineb 70 WP @25-30 g Or Azoxystrobin 18.2 % w/w + Difenconazole 11.4 % w/w SC @ 10 ml Or Fluxapyroxad 167 g/l + Pyraclostrobin 333 g/l SC @ 6g Or Carbendazim 50 WP @ 20 gm Or Propiconazole 25 EC @10 ml Or Pyraclostrobin 5% + Metiram 55% WG @ 20 g per 10 litres water. • For management of root rot, wilt, bacterial leaf blight disease in cotton it is advised for drenching at early symptomatic plants and surrounding plants with Trichoderma spp. (T. harzianum or Trichoderma viride) 1 % WP @ 50g Or Carbendazim 50 WP @ 20g/ 10 liters of water.</p>
COTTON	<p>• Considering rainfall forecast it is advised to continue the agrochemical spraying operations and inter-cultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • Spray 2 % urea at flowering stage and 2% spray of DAP at boll development stage of cotton. • Spray NAA 4.5 SL @ 4ml /10 litres of water to avoid natural shedding of squares and flowers of cotton and Mepiquat Chloride @ 10 ml/10 litres of water to restrict the excess vegetative growth of cotton after 3-4 days and current spell of rain by judging the local calm and clear weather condition. • If the incidence of thrips is noticed in cotton, on crossing ETL it is advised to spray Thiamethoxam 25% WG @ 2 gm/10L (100g/ha) Or Spinetoram 11.7 SC @ 8.4 ml/10L (420 ml/ha) after 3-4 days and current spell of rain by judging the local calm and clear weather condition. • Where the crop is at 60-90 days, if the incidence of jassids was noticed, on crossing ETL, it is advised to spray Flonicamid 50WG @ 4g/10L (200g/ha) Or Dinotefuran 20SG @ 3g/10L (150g/ha) Or Imidacloprid 17.8 SL @ 3ml/10L (150ml/ha) after 3-4 days and current spell of rain by judging the local calm and clear weather condition. • If the incidence of white fly nymph is noticed, Pyriproxyfen 10 EC @ 20ml/10L (1000 ml) /ha Or Buprofezin 25 SC @ 20ml/10L (1000 ml/ha) or Spiromesifen 22.9 SC @ 12ml/10L (600 ml/ha). • If the incidence of adult white fly is noticed, on crossing ETL spray Diafenthiuron 50% WP @ 12g/10L (600 g/ha) Or Afidopyropen 50 g/L @ 20ml/10L (1000 ml/ha) Or Dinotefuran 20 SG @ 3g/10L (150g/ha) Or Flonicamid 50 WG @4g/10L (200 g/ha) or Clothianidin 50%WDG 1ml/10L (50ml/ha) after 3-4 days and current spell of rain by judging the local calm and clear weather condition.</p>
COTTON	<p>• Considering rainfall forecast it is advised to continue the agrochemical spraying operations and inter-cultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • Monitoring of pink bollworm using pheromone traps may be initiated 45 days after sowing. Install pheromone traps @ 5 per hectare for monitoring moth activity of pink bollworm. • Inspect the crop at squaring and flowering stage of the crop for the presence of pink bollworm larvae within flowers. Remove and destroy rosette flowers whenever seen. • Where crop at below 60 days, Spray NSKE 5% + Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 1.0gm laundry detergent emulsion (Initial 1-2 sprays).</p>
Crop (Varieties)	Crop Specific Advisory
	<p>Indoxacarb 14.5 SC @ 10ml/10L (500ml/ha) Or Chlorpyrifos 20 % EC @ 25ml/10L (1250 ml/ha) after 3-4 days and current spell of rain by judging the local calm and clear weather condition. • Farmers are advised to inspect the presence and damage of pink bollworm by plucking 20 green bolls from different plants randomly (one boll per plant) if ETL have grown early damaging flower (Varieties flower advised 09 days at the crop) boll 90% bolls have out of 20 bolls. This will be helpful to increase effect on the germinal. The pheromone traps should be checked immediately in sunlight spray Prochlorvos 50 EC @ 30 ml/10L (The har/har) Or top must be harvested 50g @ 5g/10g (15th day) and is not done immediately. it should be stored at safe place protecting from rains. • If the</p>

Crop(Varieties)	<p>5 ml/ha (500 or 1500 ppm) + 1.0% laundry detergent emulsion (initial 1-2 sprays).  <b>Crop Specific Advisory</b>          NSKE 25L + Neem oil 2.5L + 0.5% Chlorpyrifos emulsion per hectare). Use          Indoxacarb 14.5% SC @ 10ml/10L (500ml/ha) Or Chlorpyrifos 20% EC @ 25ml/10L          (250 ml/ha) after 3-4 days and current spell of rain by judging the local calm and clear          weather condition.</p>
SOYABEAN	<p>pick 20 green bolls from different plants randomly (one boll per          plant) if 5% have grown &gt; 40% damaged soybean (Rosette flower) advised 09 days after the          propylthiouracil 90% past has out of 20 bolls which will not have a decrease effect on the seed          germination. The propylthiouracil should be immediately in 3 daylight spray Prochlorvos 50% EC          @ 40 ml/ha. The har/har or top must be harvest 55C @ 50 g/10g (250g/ha) and          is not done immediately, it should be stored at safe place protecting from rains. • If the          produce is to be used for seed purpose in the next season, farmers are advised to thresh          the soybean at 350 to 400 RPM thresher to avoid the loss of seed germination. • If          rainwater has accumulated in the crop area due to rains during the previous week, the          excess rainwater should be drained and also considering the forecast of the next rain,          care should be taken that the rain water does not accumulate in the crop area for a long          time. • For control the attack of Gram Pod Borers during pod filling stage, farmers are          advised to spray any one of the following insecticides: Indoxacarb 15.80 % EC (333          ml/ha), OR Flubendiamide 39.35 % w/w SC (150 ml/ha) OR Novaluron 05.25 % +          Indoxacarb 04.50 % SC (825-875 ml/ha) OR Emamectin benzoate 01.90 % EC (425          ml/ha). • If the infection of fungal diseases likes Brown Stem Rot, Pod          blight/anthracnose/Rhizoctonia Arial blight was noticed on soybean crop, farmers are          advised to apply spray of any one of the recommended fungicides like Tebuconazole          25.9 EC (625 ml/ha) OR Tebuconazole 10% + Sulphur 6 5% WG (1250 g/ha) OR          Carbendazim 12 % + Mancozeb 63 % WP (1250 g/ha) OR Picoxystrobin 22.52% w/w          SC (400 ml/ha) OR Fluxapyroxad 167 g/l + Pyraclostrobin 333 g/l SC (300 g/ha) OR          Pyraclostrobin 133 g/l + Epoxiconazole 50g/l SE (750 ml/ha) as a protective spray for          control of fungal diseases. • Farmers are advised to monitor their crop on regular          intervals preferably at 3-4 locations in their fields and see whether any          insect/pest/caterpillar is there along with their stage. This will facilitate the effective          insect control measures. • Farmers are also advised to install bird perches at different          locations which facilitate seating arrangement for predatory bird which feed on leaf          eating caterpillars. • For control of defoliators like Semilooper, tobacco caterpillar and          Heliothis, farmers are advised to spray any one of the following insecticides:          Broflanilide 300 g/l SC (42-62 g/ha) OR Flubendiamide 39.35 % w/w SC (150 ml/ha)          OR Indoxacarb 15.80 % EC (333 ml/ha), OR Tetraniliprole 18.18 SC (250-300 ml/ha)          OR Novaluron 05.25 % + Indoxacarb 04.50 % SC (825-875 ml/ha) OR          Chlorantraniliprole 18.5 % SC (150 ml/ha) OR Emamectin benzoate 01.90 % EC (425          ml/ha) OR Flubendiamide 20 % WG (250-300 g/ha) OR Lambda-cyhalothrin 04.90 %          CS (300 ml/ha) OR Profenofos 50 % EC (1 l/ha) OR Spinetoram 11.70 % SC (450          ml/ha) OR pre-mixed Beta-cyfluthrin 08.49 % + Imidacloprid 19.81 % w/w OD (350          ml/ha) OR Thiamethoxam 12.60 % + Lambda-cyhalothrin 09.50 % ZC (125 ml/ha)          OR Chlorantraniliprole 09.30 % + Lambda-cyh</p>
RICE	<p>• Considering rainfall forecast it is advised to continue the agrochemical spraying          operations and intercultural operations during next 1-2 days by taking in to account of          local clear and calm weather condition. • Apply 100 kg Nitrogen, 50 kg Phosphorus          and 50 kg Potash per hectare. Apply whole phosphorus and potash and half the amount          of nitrogen in the mud (Puddled field) and the remaining half nitrogen in two equal          installments (usually tillering stage which comes at 30 days and panicle initiation stage          which comes about 60 days). • Stem borer: - Bio-control: - Pheromone traps should be          set 20 per ha. Trichogramma japonicum (Trichocard) is a parasitic insect release          50,000 eggs per hectare 3 to 4 times every 7 days. Chemical control: - Spray          Chlorantraniliprole 0.4% G @ 10 kg. or Cartap Hydrochloride 4 G @ 18 kg. or          Fipronil 0.3 G @ 25 kg. apply per hectare when there is water in paddy bund or apply          chlorantraniliprole 18.5% SC @ 3 ml per 10 liters of water. • Gall midge: - Apply          Carbofuran 3% G @ 25 kg per hectare by maintaining water level 7 to 10 cm. Do not          remove water from paddy bunds for 4 to 5 days. These pesticides should be used again          after 30 days as required. • Blast and Neck blast: - Spray Hexaconazole 5% EC @ 20          ml. or Mencozeb 75% @ 30 gm per 10 liters of water. • Bacterial leaf blight: - Spray          Copper hydroxide 53.8% DF @ 20 gm + Streptocycline 1.5 gm per 10 liters of water.</p>
Crop(Varieties)	<p><b>Crop Specific Advisory</b>          After paddy planting till the roots of the plant are well established, the water level          should be 2-3 cm. (one inch) should be kept. After this, the level is usually about 5 cm          till the grain matures. (two inches) should be increased.</p>
CITRUS	<p>not accumulate in the crop area for a long time. • For management of Phytophthora          brown blight on fruit. Foliar spraying of Fosetyl Al for at the rate of 2.5 g per liter of          water (twice in August and September at an interval of 40 days). • Install methyl          eugenol pheromone trap at the rate of 20-25-hectare control of for fruit fly. Bait          operations and intercultural operations during next 1-2 days by taking in to account of          local clear and calm weather condition. • If rainwater has accumulated in the crop area          sucking moth at the time of colour development stage, spraying should be done of          due to rains during the previous week, the excess rainwater should be drained and also          considering the forecast of the next rain, care should be taken that the rain water does          interval, till the fruits are harvested. • The fruits fallen under the trees in the garden</p>



Crop(Varieties)	Crop Specific Advisory
CITRUS	<p>Copper hydroxide 55.8% DF @ 50 gm + Streptocycline 1.5 gm per 10 liters of water. After paddy planting till the roots of the plant are well established, the water level should not accumulate in the crop area for a long time. For management of Phytophthora blight on fruit, the water level should be kept. After this, the level is usually about 5 cm above the ground level. Foliar spraying of Fosmidil A1 fol at the rate of 2.5 g per liter of water (twice in August and September at an interval of 40 days). • Install methyl eugenol pheromone trap at the rate of 20-25-hectare control of for fruit fly. Bait operations and intercultural operations during next 1-2 days by taking in to account of (methyl eugenol) should be changed every 15-20 days. • For the management of fruit sucking moth at the time of colour development stage, spraying should be done due to rains during the previous week, the excess rainwater should be drained and also Neem oil 10 ml of petroleum spray oil (mineral oil) ml per litre of water 10-15 days considering the forecast of the next rain, care should be taken that the rain water does interval, till the fruits are harvested. • The fruits fallen under the trees in the garden should be collected and buried in the compost pit. • To increase the size of Ambia crop (June-July) fruits, spray 1.5 kg Monopotassium phosphate/ urea/Potassium nitrate/Diammonium phosphate along with Gibberlic Acid (GA<sup>3</sup>) 1.5 g in 100 litres of water. • To check the fruit dropping in Ambia Bahar a spray of Carbendazim 100 gm and GA<sup>3</sup> or 2,4-D or Naphthalene 100tic Acid (NAA) 1.5 gm in 100 liter of water.</p>

#### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
PAPAYA	<ul style="list-style-type: none"> <li>6-week-old papaya seedlings should be planted at 2.5 x 2.5 m spacing.</li> </ul>
BRINJAL	<ul style="list-style-type: none"> <li>Considering rainfall forecast it is advised to continue the agrochemical spraying operations and intercultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • Due to rainfall, higher humidity and cloudy weather condition in last week, if the incidence of fruit and shoot borer is noticed on brinjal crop, spraying should be done of any of following insecticides on crossing ETL, Carbosulfan 25 % EC 1250 ml per hectare or Deltamethrin 02.80 % EC 400 to 500 ml per hectare or Emamectin benzoate 05 % SG 200 gram per hectare or Lambda-cyhalothrin 04.90 % CS 300 ml per hectare or Spinosad 45 % SC 162 to 187 ml per hectare or Thiachloprid 21.70 % SC 750 ml per hectare or Chlorantraniliprole 09.30 % + Lambda-cyhalothrin 04.60 % ZC 200 ml per hectare mix with in 500 litres of water per hectare.</li> </ul>
TOMATO	<ul style="list-style-type: none"> <li>Considering rainfall forecast it is advised to continue the agrochemical spraying operations and intercultural operations during next 1-2 days by taking in to account of local clear and calm weather condition. • If rainwater has accumulated in the crop area due to rains during the previous week, the excess rainwater should be drained and also considering the forecast of the next rain, care should be taken that the rain water does not accumulate in the crop area for a long time. • Due to prevailing wet condition if the incidence of early blight disease is noticed in tomato crop, it is advice to take spraying during clear and calm weather condition with Azoxystrobin 23% SC 500 ml/hectare or Pyraclostrobin 20% WG @ 375-500 gm/hectare mix with in 500 litre of water per hectare on ETL.</li> </ul>

#### Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	<ul style="list-style-type: none"> <li>The floor of the animal shed should be kept dry and clean. • The feed and fodder should be stored properly to prevent the growth of moulds. • Maintain the surrounding of animal shed clean and hygienic and remove the unwanted vegetation nearby the sheds. Protect young animals from excessive consumption of newly grown green vegetation.</li> </ul>
GOAT	<ul style="list-style-type: none"> <li>Vaccinate the goat against FMD, HS, PPR and enterotoxaemia. Offer clean and cold water (stored in earthen pot) to the animals.</li> </ul>

Others (Soil / Land Preparation) Poultry(Varieties)	Others (Soil / Land Preparation) Specific Advisory Poultry Specific Advisory
BIRD	<ul style="list-style-type: none"> <li>Provide nutritious feed and fresh, clean and cold water to the birds.</li> </ul>
GENERAL ADVICE	<p>It should be noted that the validity of the dates mentioned in the Weather Based Advisory Bulletins will be valid up to 08:30 AM of the mentioned date. # Under label claim.</p>

Others (Soil / Land Preparation) Poultry (Varieties)	Others (Soil / Land Preparation) Specific Advisory Poultry Specific Advisory
BIRD  GENERAL ADVICE	<ul style="list-style-type: none"> <li>• Provide nutritious feed and fresh, clean and cold water to the birds.</li> <li>• It should be noted that the validity of the dates mentioned in the Weather Based Advisory Bulletins will be valid up to 08:30 AM of the mentioned date.</li> </ul> <p> <span style="color: blue;">#</span> Based on research that was not included in the CIBRC label claim.  <span style="color: blue;">#</span> Under label claim.         </p>

