



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Central Institute for Cotton Research,
Nagpur



Agromet Advisory Bulletin

Date : 22-02-2022

Weather Forecast of District NAGPUR(Maharashtra) Issued On : 2022-02-22(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-02-23	2022-02-24	2022-02-25	2022-02-26	2022-02-27
Rainfall(mm)	0.0	0.0	3.8	0.0	0.0
Tmax(°C)	32.8	33.6	33.1	33.8	34.2
Tmin(°C)	13.0	14.1	13.9	15.3	16.2
RH-I(%)	68	69	74	70	67
RH-II(%)	38	39	45	42	40
Wind Speed(kmph)	7.0	8.0	10.0	10.0	9.0
Wind Direction(Degree)	113	158	197	198	30
Cloud Cover(Octa)	2	2	3	2	2

Weather Summary/Alert:

In Nagpur district, as per the district level value added forecast given by, IMD, RMC, Nagpur, sky will be partly to partially cloudy weather during next five days i.e. 23rd, February, 2022 to 27th, February, 2022 and light to moderate rainfall very likely to occur at isolated places on 25th February, 2022. The maximum and minimum temperature will be 32.8 to 34.2 and 13.0 to 16.2 degree Celsius respectively and wind speed will be 7.0 to 10.0 kilometre per hour during next five days.

General Advisory:

Carry out harvesting & threshing of matured Chickpea, Wheat, Mustard, Linseed etc. and store the harvested produce in safer places, if threshing is not possible cover the harvested produce with plastic sheet. It is advised to continue the intercultural operation, spraying of agrochemicals for pest and disease management by considering the soil moisture and crop need in rabi season crop, fruit orchards and vegetables by judging local clear weather condition.

SMS Advisory:

Carry out harvesting & threshing of matured Chickpea, Wheat, Mustard, Linseed etc. and store the harvested produce in safer places, if threshing is not possible cover the harvested produce with plastic sheet.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BENGAL GRAM/ CHICK PEA	Carry out harvesting & threshing of matured chickpea and store the harvested produce in safer places, if threshing is not possible cover the harvested produce with plastic sheet. In late sown and irrigated chickpea if Gram Pod Borer has reached the ETL, spraying should be done with Quinalphos 25% EC 20 ml or Emamectin benzoate 5% SG 3 gm or Flubendiamide 20% WG 5 gm or Chlorantraniliprole 18.5 SC 2.5 ml per 10 liters of water.

Crop(Varieties)	Crop Specific Advisory
WHEAT	Carry out harvesting & threshing of matured wheat and store the harvested produce in safer places, if threshing is not possible cover the harvested produce with plastic sheet. Considering the soil moisture status and crop needs, wheat crop should be irrigated at flowering stage (65 to 70 days after sowing) and milking stage (80 to 85 days after sowing). In wheat, in case of infestation of rust and leaf blight disease is noticed spraying should be done of Mancozeb 75 % WP @ 20 to 25 gm per 10 liters of water.
GROUNDNUT	In summer groundnut, gap filling should be done up to 10 days after emergence

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
MANGO	If the infestation of jassids on mango is crossing economic threshold level, Buprofezin 25 % EC 10 ml or deltamethrin 2.8 % EC 5 ml or imidacloprid 17.8 SL % 4 ml or Lambda cyhalothrin 5 % EC 5 ml or Dimethoate 30% EC 16.5 ml per 10 liters of water. For management of powdery mildew of mango, spraying should be done of Hexaconazole 5 % EC 10 ml or Penconazole 10 % EC 5 ml or Sulfur 80 % WDG 30 gm or Dinocap 48 % EC 5 ml per 10 liters of water.
ORANGE	During this period in general citrus leaf miner and citrus psylla are active. At the time of Ambia flush for citrus psylla control, spray imidacloprid 0.5 ml, quinalphos 2 ml or dimethoate 2 ml in one litre water. Against leaf miner spray thiamethoxam 0.3 g or Fenvalerate 1 ml per litre of water. Repeat application after 10 days in case the infestation of leaf miner and citrus psylla persists. In case of citrus aphid attack, spray quinalphos @ 2 ml or dimethoate @ 2 ml/l and repeat it after every 7 days. In case there is gummosis on the tree trunk remove the gum portion by sharp knife and apply Metalaxyl paste or fosetyl Al paste on the affected part of the tree trunk. During this period incidence of lemon butterfly is observed. Spray Quinalphos 2 ml or Fenvalerate 2 ml in one liter water for minimizing the incidence of lemon butterfly. Foliar application of gibberellic acid 1.5 gm + urea 1 kg may be given to Ambia bahar during fruit set after mixing it in 100 litres water. (Dissolve gibberellic acid in 10 ml alcohol before mixing in water)
CHILLI	In chilli, if thrips, fruit borer and tobacco leaf eating caterpillar crosses ETL spraying should be done of Cyantraniliprole 10.26 OD at the rate of 12.0 ml per 10 litres of water during clear weather condition.
CABBAGE	In cabbage, if aphids (green & black), diamond back moth and tobacco leaf eating caterpillar crosses ETL spraying should be done of Cyantraniliprole 10.26 OD at the rate of 12.0 ml per 10 litres of water during clear weather condition.
TOMATO	In tomato, if leaf minor, aphids, thrips, white fly and fruit borer crosses ETL spraying should be done of Cyantraniliprole 10.26 OD at the rate of 18.0 ml per 10 litres of water during clear weather condition.
LINSEED	Carry out harvesting & threshing of matured Linseed and store the harvested produce in safer places, if threshing is not possible cover the harvested produce with plastic sheet. In linseed, if the gall midge crosses economic threshold level (8.77 % gall midge) spraying should be done of Acetamiprid 20 SP 2.0 gram per 10 litres of water or Imidacloprid 17.8 SL 2.5 ml per 10 litres of water or Dimethoate 30 % EC 10 ml per 10 liters of water. Mature linseed crop should be harvested and stored in a safe place.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	In case of infestation of ticks and fleas on the body of goats and goats' kids, rub the goats and kids with equal quantity of ash and 50 % Carbaryl powder. In case of high incidence, add 2 ml of Butox (Deltamethrin 12.5 mg) or Amitraz and bathe the goats and kids. Pesticides should be used very carefully and care should be taken not to poison the goats by licking them. Goats and sheep should be vaccinated against rinder pest disease.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Considering the availability of irrigation and market demand in summer season, leafy vegetables, okra, guar, chawli, cucumber, Bottle Gourd, pumpkin, ridge gourd, bitter gourd, round gourd, watermelon, muskmelon, spinach, radish, fenugreek and coriander should be cultivated.
GENERAL ADVICE	It should be noted that the validity of the dates mentioned in the Weather Based Advisory Bulletins will be valid from 08:30 AM on the previous day to 08:30 AM on the said date.